International Application No.: PCT/DE2003/002030

International Filing Date: June 18, 2003

Page: 5

Amendments to the Claims

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

- (Currently Amended) A method for producing reactive organic compounds containing poly-DOPO, said method comprising which are obtained by addition of reacting 9,10-dihydro-9-oxa-10-phosphaphenanthrene-10-oxide (DOPO) onto with acetylenically unsaturated compounds which carry having reactive groups, wherein said reacting step is performed in the presence of a catalyst which is suitable for the addition of triple bonds.
- 2. (Currently Amended) The method as claimed in claim 1, wherein the organophosphorus compound DOPO and the acetylenically unsaturated compound are reacted with one another in the a ratio of 1.5 to 3 mol of DOPO per triple bond[[,]] preferably 1.9 to 2.1 mol of DOPO per triple bond.
- 3. (Currently Amended) The method as claimed in claim 1 and 2, wherein the acetylenically unsaturated compounds used are alkynes, alkynols, alkynecarboxylic acids, alkynecarboxylic esters or corresponding alkadiyne compounds.
- 4. (Currently Amended) The method as claimed in claim 1 to 3, wherein the catalyst used for the reaction of the organophosphorus compound DOPO with the acetylenically unsaturated compound is mercury salts, or copper salts, or amines or [[,]] preferably [[,]] aluminum triisopropoxide.
- 5. (Currently Amended) The method as claimed in claim 1 to 4, wherein the addition reaction is carried out in solution[[,]] where the solvent used is preferably 1,4 dioxane.

International Application No.: PCT/DE2003/002030

International Filing Date: June 18, 2003

Page: 6

6. (Currently Amended) Flame retardant for thermoplastic polymers comprising The use of the reactive organic compounds containing poly-DOPO prepared as in claims 1 to 5 as flame retardant for thermoplastic polymers.

- 7. (New) The method as claimed in Claim 2, wherein the ratio is 1.9 to 2.1 mol of DOPO per triple bond.
- 8. (New) The method as claimed in Claim 5, wherein the addition reaction is carried out in 1,4-dioxane.